

doi:10.1093/cvr/cvv051

Published online 27 February 2015

Status of P2Y₁₂ treatment must be considered in evaluation of myocardial ischaemia/reperfusion injury

We noted with interest the recent publication by the ESC Working Group on pre-clinical assessment of novel cardioprotective therapies.¹ Because of the nearly three decades of experimental work which has not yet yielded a single intervention that has been demonstrated to diminish the amount of infarcting myocardium following ischaemia/reperfusion in patients, it is necessary to critically examine the available data to determine where there might be roadblocks. Lecour *et al.* have analysed many possible confounders that might be responsible for the failure to extrapolate observations in the animal laboratory to the cardiac catheterization suite. They are to be congratulated for this succinct summary. However, we are disappointed that they have overlooked a particularly compelling explanation for the failures. Patients with acute myocardial infarction are treated with numerous pharmacological agents before the actual percutaneous coronary intervention that will open the coronary artery. One of these agents is clopidogrel, a platelet P2Y₁₂ receptor antagonist, which for the past decade has been administered to all patients according to published guidelines. We have shown that it and other P2Y₁₂ antagonists have cardioprotective properties which are independent of their anti-thrombotic actions.^{2–6} We suspect that this anti-infarct effect may well be responsible for their resounding clinical success. We suggest that any other intervention applied in such patients which uses a similar mechanism of protection as the administered P2Y₁₂ antagonist will not provide additional protection. Indeed, neither ischaemic pre-² nor post-conditioning⁶ caused any additional protection in our animal models treated with a maximally protective dose of a P2Y₁₂ inhibitor. Only an intervention that uses a different mechanism of protection can be expected to provide incremental protection. We all agree that additional protection is needed for these patients and that the search for cardioprotective interventions must continue. However, we submit P2Y₁₂ inhibitors constitute an important confounder in the clinical studies, and strongly suggest that testing of future candidates for cardioprotection be examined in animal models also treated with a P2Y₁₂ agent⁶ before submitting the new intervention to costly clinical trials.

References

- Lecour S, Bøtker HE, Condorelli G, Davidson SM, Garcia-Dorado D, Engel FB, Ferdinandy P, Heusch G, Madonna R, Ovize M, Ruiz-Meana M, Schulz R, Sluijter JPG, Van Laake LW, Yellon DM, Hausenloy DJ. ESC Working Group Cellular Biology of the Heart: position paper: improving the preclinical assessment of novel cardioprotective therapies. *Cardiovasc Res* 2014;**104**: 399–411.
- Yang X-M, Liu Y, Cui L, Yang X, Liu Y, Tandon N, Kambayashi J, Downey JM, Cohen MV. Platelet P2Y₁₂ blockers confer direct postconditioning-like protection in reperfused rabbit hearts. *J Cardiovasc Pharmacol Ther* 2013;**18**:251–262.
- Yang X-M, Liu Y, Cui L, Yang X, Liu Y, Tandon N, Kambayashi J, Downey JM, Cohen MV. Two classes of anti-platelet drugs reduce anatomical infarct size in monkey hearts. *Cardiovasc Drugs Ther* 2013;**27**:109–115.
- Yang X-M, Cui L, Alhammouri A, Downey JM, Cohen MV. Triple therapy increased myocardial salvage during ischemia/reperfusion in the in situ rat heart. *Cardiovasc Drugs Ther* 2013;**27**:403–412.
- Cohen MV, Downey JM. Combined cardioprotectant and antithrombotic actions of platelet P2Y₁₂ receptor antagonists in acute coronary syndrome: just what the doctor ordered. *J Cardiovasc Pharmacol Ther* 2014;**19**: 179–190.
- Cohen MV, Downey JM. Signalling pathways and mechanisms of protection in pre- and postconditioning: historical perspective and lessons for the future. *Br J Pharmacol* 2015; in press.

Michael V. Cohen^{1,*} and James M. Downey²

¹Department of Physiology and Cell Biology and Medicine, University of South Alabama College of Medicine, Mobile, AL, USA

²Department of Physiology, Emeritus, University of South Alabama College of Medicine, Mobile, AL, USA

*Corresponding author. Tel: +1 251 460 6812; fax: +1 251 460 6464. Email: mcohen@southalabama.edu

doi:10.1093/cvr/cvv052

Published online 27 February 2015

Platelet inhibitors influence cardioprotection: importance in preclinical study design: reply

We would like to thank Professors Cohen and Downey for highlighting platelet inhibitors, in particular platelet P2Y₁₂ receptor antagonists, as an important confounder to take into consideration in pre-clinical studies designed to study novel cardioprotective strategies against ischaemia/reperfusion injury. As already mentioned in our recent publication from the ESC working group, both clinical and animal studies give evidence that platelet inhibitors reduce myocardial infarct size by mechanisms that may involve nitric oxide or adenosine.^{1–3} Over the past few years, Cohen and Downey have published strong and

convincing evidence suggesting that additional protection cannot be afforded with pharmacological and/or mechanical protective strategies sharing the same pro-survival signalling pathways as the P2Y₁₂ antagonist (see review³). Therefore, we fully share their point of view that the effect of co-medication with P2Y₁₂ antagonists or with any other medications frequently given to ischaemic heart disease patients (e.g. aspirin, statins, ACE inhibitors, beta-blockers, etc.)¹ on cardioprotective therapies needs to be tested in pre-clinical settings prior to translation to clinical cardioprotection as also recently reviewed in detail by Ferdinandy *et al.*⁴

References

- Lecour S, Bøtker HE, Condorelli G, Davidson SM, Garcia-Dorado D, Engel FB, Ferdinandy P, Heusch G, Madonna R, Ovize M, Ruiz-Meana M, Schulz R, Sluijter JPG, Van Laake LW, Yellon DM, Hausenloy DJ. ESC Working Group Cellular Biology of the Heart: Position paper: improving the preclinical assessment of novel cardioprotective therapies. *Cardiovasc Res* 2014;**104**: 399–411.
- Roubille F, Lairez O, Mewton N, Rioufol G, Ranc S, Sanchez I, Cung TT, Elbaz M, Piot C, Ovize M. Cardioprotection by clopidogrel in acute ST-elevated myocardial infarction patients: a retrospective analysis. *Basic Res Cardiol* 2012;**107**:275.
- Cohen MV, Downey JM. Combined cardioprotectant and antithrombotic actions of platelet P2Y₁₂ receptor antagonists in acute coronary syndrome: just what the doctor ordered. *J Cardiovasc Pharmacol Ther* 2014;**19**: 179–190.
- Ferdinandy P, Hausenloy DH, Heusch G, Baxter GF, Schulz R. Interaction of risk factors, comorbidities and comedications with ischemia/reperfusion injury and cardioprotection by preconditioning, postconditioning, and remote conditioning. *Pharmacol Rev* 2014;**66**: 1142–1174.

Sandrine Lecour¹, Rainer Schulz², Péter Ferdinandy^{3,4} and Derek J. Hausenloy^{5,6,7,8,*}

¹Department of Medicine, Hatter Institute for Cardiovascular Research in Africa and MRC Inter-University Cape Heart Group, University of Cape Town, Cape Town, South Africa

²Institute of Physiology, Justus-Liebig University, Giessen, Germany

³Clinical Pharmacology, Department of Pharmacology and Pharmacotherapy, Semmelweis University, Budapest, Hungary

⁴Pharmahungary Group, Szeged, Hungary

⁵The Hatter Cardiovascular Institute, Institute of Cardiovascular Science, University College London, London, UK

⁶The National Institute of Health Research University College London Hospitals Biomedical Research Centre, London, UK

⁷National Heart Research Institute Singapore, National Heart Centre Singapore, Singapore

⁸Cardiovascular and Metabolic Disorders Program, Duke-National University of Singapore, Singapore

*Corresponding author. Email: d.hausenloy@ucl.ac.uk